arXiv:1207.1252v2 [math-ph] 2 Aug 2012

the Dimer Gas Mayer Series, the Monomer-Dimer $\lambda_d(p)$, the Federbush Relation

Paul Federbush
Department of Mathematics
University of Michigan
Ann Arbor, MI 48109-1043
(`pfed@umich.edu`)

August 3, 2012

Abstract

The author and Shmuel Friedland recently presented an expression for $\lambda_d(p)$ of the monomer-dimer problem involving a power series in p . Herein I present a simple way to derive this expression for $\lambda_d(p)$ from the Mayer (or Virial) series of a dimer gas. The derivation is an exercise in basic statistical mechanics. We expect the relationship to be very useful.

Shmuel Friedland and the author recently presented an expression for $\lambda_d(p)$ of the monomer-dimer problem involving a power series in p, see eq (7.1) of [1]. We always knew that the Mayer or Virial series of a dimer gas implicitly contained all the information to compute our series. (In [2], see the remark there in the paragraph following the paragraph containing eq. (30), the sentence beginning: "By a simple device...".) But it was not until the referee of a recent paper, [3], asked about the relationship of the expansion for $\lambda_d(p)$ to the Virial expansion that I finally worked out the details of the simple natural connection. We present the general development, and then specialize to $d = 2$ where the most is now known.

We take the statistical mechanical notation from Ruelle, [4]. We must be careful to avoid confusions, for example between the p in $\lambda_d(p)$ and pressure. We at the outset specify some notation.

1) p (as in $\lambda_d(p)$) is considered the <u>density</u> of vertices covered by dimers.

2) ρ is the <u>density</u> of dimers. So

$$p = 2\rho. \tag{1}$$

3) P is <u>pressure</u>.

4) The <u>Mayer series</u> from [4], page 84 are

$$\beta P = \sum_{n=1}^{\infty} b_n z^n \tag{2}$$

and

$$\rho = \sum_{n=1}^{\infty} n b_n z^n. \tag{3}$$

We will set $\beta = 1$. But we choose to work with the density p so we replace (3) by

$$p = 2 \sum_{1}^{\infty} n b_n z^n. \tag{4}$$

5) We solve (4) for $z = z(p)$ by iterating (from $z = 0$)

$$z = \frac{1}{2b_1} p - \sum_{2}^{\infty} \frac{n b_n}{b_1} z^n. \tag{5}$$

We then substitute $z = z(p)$ into the right side of (2), getting

$$P(p) = \sum_{1}^{\infty} b_n (z(p))^n \tag{6}$$

Written as a power series in p this is the Virial series.

Considering the partition function leading to the Mayer series, we note that in volume V the sum is dominated by terms with $\sim \rho V$ dimers, leading to the observation that

$$\lambda_d(p) = P(p) - \frac{p}{2} \ln(z(p)) \tag{7}$$

This is the Federbush relation. (But perhaps this equality already has some name?) We here assume that the expansion of the left side of this relation given in eq. (7.1) of [1] is consistent with this relation, 'as it certainly must be'. We discuss this at the end of the paper.

We look at $z(p)$ as derived above

$$z = \frac{p}{2b_1}(1 + F(p)) \tag{8}$$

where $F(p)$ is a power series in p without a constant term.We use (8) to get

$$\lambda_d(p) = P(p) - \frac{p}{2} \ln\left(\frac{p}{2b_1}\right) - \frac{p}{2} \ln(1 + F(p)) \tag{9}$$

$$= -\frac{p}{2} \ln(p) + \frac{p}{2} \ln(2d) + P(p) - \frac{p}{2} \ln(1 + F(p)) \tag{10}$$

We have used that $b_1 = d$. Aside from the first two terms with ln's, the last two terms in (10) are polynomial series in p.

For $d = 2$ we calculated as the principal part of our computer computations (in calculating the $\bar{J}_i$ defined in [1]) the values of the first 7 coefficients in the Mayer series of a dimer gas:

$$b_1 = 2 \tag{11}$$

$$b_2 = -7 \tag{12}$$

$$b_3 = \frac{116}{3} \tag{13}$$

$$b_4 = -\frac{521}{2} \tag{14}$$

$$b_5 = \frac{9812}{5} \tag{15}$$

$$b_6 = -\frac{47644}{3} \tag{16}$$

$$b_7 = \frac{945688}{7} \tag{17}$$

Equation (10) with the appropriate substitutions yields the same series as in [3], up to order p^7 in the power series. The computations are easy using Maple:

$$\begin{aligned}\lambda_2(p) \sim &\frac{1}{2}(p\ln(4) - p\ln p - 2(1-p)\ln(1-p) - p) + \frac{4}{2}\cdot\left(\frac{1}{2\cdot 1}\left(\frac{p}{4}\right)^2\right. \\ &\left. + \frac{1}{3\cdot 2}\left(\frac{p}{4}\right)^3 + \frac{7}{4\cdot 3}\left(\frac{p}{4}\right)^3 + \frac{41}{5\cdot 4}\left(\frac{p}{4}\right)^5 + \frac{181}{6\cdot 5}\left(\frac{p}{4}\right)^6 + \frac{757}{7\cdot 6}\left(\frac{p}{4}\right)^7\right)\end{aligned} \tag{18}$$

We previously proved that the series for $\lambda_d(p)$ in [1] converged for small p, [5],but we did not prove it converged to the correct value. The series in (10) clearly converges to the correct value. At first we felt one should be able to identify the two series just by algebraic manipulation. This is clearly possible up to any given fixed power of p. But we are now of the opinion that only a proof using difficult complex-analytic techniques offers itself. We hope to accomplish such a proof in the coming year(s).

As we stated above the two series are equal to power p^7, using the values of the b_i in (11) - (17). We made an "arbitrary" fixed integer-valued choice for the b_i for $i > 7$ and verified the equality of the two series up to power p^{33}, for this choice. This is pretty convincing proof that the two series are equal.

References

[1] P. Federbush and S. Friedland, "An Asymptotic Expansion and Recursive Inequalities for the Monomer-Dimer Problem," *Jour. of Stat. Phys* 143 (2011), 306.

[2] P. Federbush, "Computation of Terms in the Asymptotic Expansion of Dimer λ_d for High Dimensions," *Phys. Lett. A* 374 (2009), 131-133.

[3] P. Federbush, "Asymptotic Expansions for λ_d of the Dimer and Monomer-Dimer Problems," arXiv:1206.6241

[4] Ruelle, David, *Statistical Mechanics*, W. A. Benjamin, Inc. Amsterdam, 1969

[5] P. Federbush, "Convergence of the Formal Expansion for $\lambda_d(p)$ of the Monomer-Dimer Problem for Small p," arXiv:1101.4591.